

The inverse Sturm-Liouville problem with mixed boundary conditions

Dmitri Chelkak* and Evgeny Korotyaev†

February 2, 2008

Abstract

Consider the operator $H\psi = -\psi'' + q\psi = \lambda\psi$, $\psi(0) = 0$, $\psi'(1) + b\psi(1) = 0$ acting in $L^2(0, 1)$, where $q \in L^2(0, 1)$ is a real potential. Let $\lambda_n(q, b)$, $n \geq 0$, be the eigenvalues of H and $\nu_n(q, b)$ be the so-called norming constants. We give a complete characterization of all spectral data $(\{\lambda_n\}_0^\infty; \{\nu_n\}_0^\infty)$ that correspond to $(q; b) \in L^2(0, 1) \times \mathbb{R}$. If b is fixed, then we obtain a similar characterization and parameterize the iso-spectral manifolds.

1 Introduction and main results

Consider the Sturm-Liouville problem

$$-\psi'' + q(x)\psi = \lambda\psi, \quad x \in [0, 1], \quad q \in L^2(0, 1),$$

with boundary conditions

$$\psi'(0) - a\psi(0) = 0, \quad \psi'(1) + b\psi(1) = 0, \quad a, b \in \mathbb{R} \cup \{\infty\},$$

where real q belongs to $L^2(0, 1)$, equipped with the norm $\|q\|^2 = \int_0^1 q(t)^2 dt$. There are a lot of papers (Borg, Gel'fand, Levitan, Marchenko, Trubowitz, ...) devoted to the inverse spectral theory for Sturm-Liouville problems on a finite interval (see the books [L], [M], [PT]). Recall that the inverse problem consists of the following parts:

- i) Prove that the spectral data (eigenvalues and some "additional" parameters) **uniquely** determine the potential.
- ii) **Reconstruct** the potential from spectral data.
- iii) **Characterize** all spectral data that correspond to some fixed class of potentials.

*Correspondence author. Dept. of Math. Analysis, Math. Mech. Faculty, St.Petersburg State University. Universitetskij pr. 28, Staryj Petergof, 198504 St.Petersburg, Russia, e-mail: delta4@math.spbu.ru

†Institut für Mathematik, Humboldt Universität zu Berlin, Rudower Chaussee 25, 12489, Berlin, Germany, e-mail: evgeny@math.hu-berlin.de

The first uniqueness results were obtained by Borg. The first results about the reconstruction were obtained by Gel'fand and Levitan (see details in [L], [M]). Trubowitz and his co-authors ([IT], [IMT], [DT], [PT]) developed the "analytic" method based on nonlinear functional analysis and the explicit reconstruction procedure for the special case when only *one* spectral datum has been changed. The book [PT] contains an elegant complete solution of the inverse Dirichlet (i.e. $a = b = \infty$) problem on $[0, 1]$. In [IT] the case when (non fixed) a, b run through $\mathbb{R}$ was considered (we recall the main result of [IT] in Theorem 5.1) and [IMT] is devoted to the case of fixed $a, b \in \mathbb{R}$. These authors ([IT], [IMT]) reduce the mapping $(q, a, b) \mapsto \{\text{spectral data}\}$ to the case of Dirichlet boundary conditions [PT]. This reduction allows to prove the characterization theorem, to construct the isospectral manifolds (i.e. the set of all (q, a, b) with the same spectrum) and so on.

Consider the case of mixed boundary condition $a = \infty$, $b \in \mathbb{R}$. We think that the best results in this case are those of Dahlberg and Trubowitz [DT]. In order to describe their results we need some definitions. Define the self-adjoint operator H in $L^2(0, 1)$ by

$$H\psi = -\psi'' + q(x)\psi, \quad \psi(0) = 0, \quad \psi'(1) + b\psi(1) = 0, \quad b \in \mathbb{R}.$$

Here and below $(') = \frac{\partial}{\partial x}, (\cdot) = \frac{\partial}{\partial \lambda}$. Denote by $\{\lambda_n(q, b)\}_0^\infty$ the eigenvalues of H . It is well-known that all $\lambda_n(q, b)$ are simple and satisfy the asymptotics

$$\lambda_n(q, b) = \lambda_n^0 + Q_0 + 2b + \mu_n(q, b), \quad \text{where} \quad Q_0 = \int_0^1 q(t)dt \quad \text{and} \quad \{\mu_n(q, b)\}_0^\infty \in \ell^2 = \ell_0^2.$$

Here and below the real Hilbert spaces ℓ_m^2 are given by $\ell_m^2 = \{\{f_n\}_0^\infty : \sum_{n \geq 0} (n+1)^{2m} f_n^2 < \infty\}$, $m \geq 0$, and $\lambda_n^0 = \pi^2(n + \frac{1}{2})^2$, $n \geq 0$, are the unperturbed eigenvalues. The monotonicity property $\lambda_0 < \lambda_1 < \dots$ gives that if $(q; b)$ runs through $L^2(0, 1) \times \mathbb{R}$, then $\{\mu_n\}_0^\infty$ doesn't run through the whole space ℓ^2 . In order to describe this situation, we introduce the open and convex set

$$\mathcal{M} = \{\{\mu_n\}_0^\infty \in \ell^2 : \lambda_0^0 + \mu_0 < \lambda_1^0 + \mu_1 < \dots\} \subset \ell^2.$$

Fix some $b \in \mathbb{R}$. The main results of [DT] are:

- (i) *The sequence of real numbers $\lambda_n^* = \lambda_n^0 + c^* + \mu_n^*$, $c^* \in \mathbb{R}$, is the spectrum of H for some potential $q \in L^2(0, 1)$ if and only if $\{\mu_n^*\}_0^\infty \in \mathcal{M}$.*
- (ii) *Let $q \in L^2(0, 1)$ and $e \in L_{\text{even}}^2(0, 1)$, i.e. $e(1-x) = e(x)$, $x \in [0, 1]$. There are two cases: (a) there is no potential $p \in L^2(0, 1)$ such that the Dirichlet spectrum of p coincides with the Dirichlet spectrum of e and $\lambda_n(p, b) = \lambda_n(q, b)$ for all $n \geq 0$; (b) such a potential p is unique. An explicit condition which distinguishes the cases (a) and (b) from each other is given in terms of $\{\lambda_n(q, b)\}_0^\infty$ and the Dirichlet spectrum of e .*

Roughly speaking, (ii) describes the bijection between the set of isospectral potentials

$$\text{Iso}_b(\{\lambda_n^*\}_0^\infty) = \{q \in L^2(0, 1) : \lambda_n(q, b) = \lambda_n^* \text{ for all } n \geq 0\}. \quad (1.1)$$

and some open subset of $L_{\text{even}}^2(0, 1)$ (see [DT] for details).

The main goal of our paper is to give a more explicit characterization of spectral data in the style of [PT], [IT] and to parameterize the isospectral manifolds in a more classic way.

Let $\varphi(x) = \varphi(x, \lambda, q)$ and $\xi_b = \xi_b(x, \lambda, q)$ be solutions of $-\psi'' + q(x)\psi = \lambda\psi$ such that

$$\varphi(0) = 0, \quad \varphi'(0) = 1 \quad \text{and} \quad \xi_b(1) = -1, \quad \xi_b'(1) = b.$$

Note that the eigenvalues $\lambda_n(q, b)$ are the roots of the Wronskian

$$w(\lambda) = w(\lambda, q, b) = \{\varphi, \xi_b\}(\lambda, q) = \varphi'(1, \lambda, q) + b\varphi(1, \lambda, q),$$

where $\{\varphi, \xi_b\} = \varphi\xi_b' - \varphi'\xi_b$. The Hadamard Factorization Theorem implies the identity

$$w(\lambda, q, b) = \cos \sqrt{\lambda} \cdot \prod \frac{\lambda - \lambda_n(q, b)}{\lambda - \lambda_n^0}, \quad \lambda \in \mathbb{C}. \quad (1.2)$$

Let $\psi_n(x) = \psi_n(x, q, b)$ be the n -th normalized eigenfunction of H such that $\psi_n'(0) > 0$. We introduce the norming constants ("additional" spectral data) by

$$\nu_n(q, b) = \log [(-1)^n \varphi(1, \lambda_n(q, b), q)] = \log \left| \frac{\psi_n(1, q, b)}{\psi_n'(0, q, b)} \right|, \quad n \geq 0, \quad (1.3)$$

$$\nu_n^0 = \nu_n(0, 0) = -\log k_n^0, \quad \text{where} \quad k_n^0 = \sqrt{\lambda_n^0} = \pi(n + \tfrac{1}{2}).$$

Recall that $\lambda_n(q, b) = \lambda_n^0 + Q_0 + 2b + \mu_n(q, b)$. Our main result is

Theorem 1.1. (i) *The mapping*

$$\Phi : (q; b) \mapsto (Q_0 + 2b; \{\mu_n(q, b)\}_0^\infty; \{\nu_n(q, b) - \nu_n^0\}_0^\infty)$$

is a real-analytic isomorphism between $L^2(0, 1) \times \mathbb{R}$ and $\mathbb{R} \times \mathcal{M} \times \ell_1^2$.

(ii) *For each $(q; b) \in L^2(0, 1) \times \mathbb{R}$ the following identity is fulfilled:*

$$b = \sum_{n \geq 0} \left(2 - \frac{e^{\nu_n(q, b)}}{|\dot{w}(\lambda_n, q, b)|} \right). \quad (1.4)$$

Remarks. i) In the proof of (i) we use the method from [PT]. The main ingredients are nonlinear functional analysis and the explicit reconstruction procedure, when only one λ_n or ν_n has been changed. Using similar arguments, it is possible to reprove the main result of [IT], i.e. the complete characterization of spectral data in the case $(q, a, b) \in L^2(0, 1) \times \mathbb{R}^2$, without the reduction to the inverse Dirichlet problem.

ii) Identity (1.4) gives the explicit expression of b in terms of spectral data (in the case $a = \infty$). This can be rewritten in the form $b = \sum_{n \geq 0} (2 - \|\xi_b(\cdot, \lambda_n, q)\|_{L^2(0, 1)}^{-2})$ (see Lemma 2.2). Note that similar identities were proved in [JL] using the technique of transmutation operators and the Gel'fand-Levitan equation. Our proof is based on the contour integration. In Sect. 5 (Appendix) we prove the analogue of (1.4) in the case $a, b \in \mathbb{R}$.

Fix some $b \in \mathbb{R}$ (for instance, $b=0$ gives the boundary conditions $\psi(0)=0, \psi'(1)=0$). In this case spectral data $\{\lambda_n\}_0^\infty, \{\nu_n\}_0^\infty$ are not independent since they satisfy the nonlinear equation (1.4). Fortunately, the first eigenvalue $\lambda_0(q, b)$ can be uniquely reconstructed from the other spectral data (i.e. $\{\lambda_n\}_1^\infty$ and $\{\nu_n\}_0^\infty$). More precisely, we have

Corollary 1.2. *For any fixed $b \in \mathbb{R}$ the mapping*

$$\Phi_b : q \mapsto (Q_0; \{\mu_{n+1}(q, b)\}_0^\infty, \{\nu_n(q, b) - \nu_n^0\}_0^\infty)$$

is a real-analytic isomorphism between $L^2(0, 1)$ and $\mathbb{R} \times \mathcal{M}^{(1)} \times \ell_1^2$, where

$$\mathcal{M}^{(1)} = \{\{\mu_{n+1}\}_0^\infty \in \ell^2 : \lambda_1^0 + \mu_1 < \lambda_2^0 + \mu_2 < \dots\} \subset \ell^2.$$

It is possible to "remove" from the spectral data not only the first eigenvalue λ_0 but also one norming constant. Recall that for any sequence $\{\lambda_n^*\}_0^\infty$ such that $\lambda_n^* = \lambda_n^0 + c^* + \mu_n^*$, where $(c^*; \{\mu_n^*\}_0^\infty) \in \mathbb{R} \times \mathcal{M}$, the set of isospectral potentials $\text{Iso}_b(\{\lambda_n^*\}_0^\infty)$ is defined by (1.1). Note that

$$w(\lambda, q, b) = w^*(\lambda) = \cos \sqrt{\lambda} \cdot \prod_{n \geq 0} \frac{\lambda - \lambda_n^*}{\lambda - \lambda_n^0} \quad \text{for each } q \in \text{Iso}_b(\{\lambda_n^*\}_0^\infty).$$

Corollary 1.3. *Fix some $b \in \mathbb{R}$ and let $(c^*; \{\mu_n^*\}_0^\infty) \in \mathbb{R} \times \mathcal{M}$. Then for each $m \geq 0$ the mapping*

$$q \mapsto \{\nu_n(q, b) - \nu_n^0\}_{n=0, n \neq m}^\infty$$

is a real-analytic isomorphism between $\text{Iso}_b(\{\lambda_n^*\}_0^\infty)$ and the open nonempty set $\mathcal{N}_m^b \subset \ell_1^2$ given by

$$\mathcal{N}_m^b = \left\{ \{\nu_n - \nu_n^0\}_{n=0, n \neq m}^\infty \in \ell_1^2 : \sum_{n: n \neq m} \left(2 - \frac{e^{\nu_n}}{|\dot{w}^*(\lambda_n^*)|} \right) > b - 2 \right\}. \quad (1.5)$$

Remark. Let $\{\nu_n - \nu_n^0\}_{n: n \neq m} = t\gamma$, where $t \in \mathbb{R}$, $\gamma = \{\gamma_n\}_{n: n \neq m} \in \ell_1^2$ and $\gamma_n > 0$, $n \neq m$. Then, due to the monotonicity reasons, there exists $t_0 \in \mathbb{R}$ such that $t\gamma \in \mathcal{N}_m^b$ iff $t < t_0$.

In conclusion, note that the method from [PT] works well in other inverse spectral problems with a purely discrete spectrum. In particular, this scheme was applied in the paper [CKK] devoted to the inverse problem for the perturbed harmonic oscillator on $\mathbb{R}$. The inverse problem for the harmonic oscillator in $\mathbb{R}^3$ perturbed by a spherically symmetric potential is reformulated as a problem on the half-line $\mathbb{R}_+$ and it is solved in [CK]. In the last case the boundary conditions are essentially nonsymmetric and it gives an extra motivation to investigate the case of mixed conditions on $[0, 1]$, which are also essentially nonsymmetric.

We describe the plan of the paper. In Sect. 2 we prove identity (1.4) and the Uniqueness Theorem. Sect. 3 is devoted to the analytic properties of the mapping Φ . In Sect. 4 we prove our main results: Theorem 1.1 and Corollaries 1.3, 1.2. Sect. 5 (Appendix) is devoted to the modifications of identity (1.4) and Corollary 1.2 to the case $a, b \in \mathbb{R}$.

Acknowledgments. Dmitry Chelkak was partly supported by grants VNP Minobrazovaniya 3.1–4733, RFFR 03–01–00377 and NSh–2266.2003.1. Evgeny Korotyaev was partly supported by DFG project BR691/23-1. Some part of this paper was written at the Mittag-Leffler Institute, Stockholm. The authors are grateful to the Institute for the hospitality. The authors would like to thank Markus Klein for useful discussions.

2 Preliminaries and proof of identity (1.4)

Introduce the fundamental solutions $\vartheta(x, \lambda, q)$, $\varphi(x, \lambda, q)$ of the differential equation

$$-\psi'' + q(x)\psi = \lambda\psi, \quad \lambda \in \mathbb{C}, \quad (2.1)$$

such that

$$\vartheta(0, \lambda, q) = \varphi'(0, \lambda, q) = 1, \quad \vartheta'(0, \lambda, q) = \varphi(0, \lambda, q) = 1.$$

For each $x \in [0, 1]$ the functions $\vartheta, \vartheta', \varphi, \varphi'$ are entire with respect to $(\lambda, q) \in \mathbb{C} \times L^2_{\mathbb{C}}(0, 1)$ (see [PT]). Moreover, the following asymptotics are fulfilled:

$$\vartheta(x, \lambda, q) = \cos \sqrt{\lambda}x + \frac{1}{2\sqrt{\lambda}} \int_0^x \left(\sin \sqrt{\lambda}x + \sin \sqrt{\lambda}(x-2t) \right) q(t)dt + O\left(\frac{e^{|\operatorname{Im} \sqrt{\lambda}|x}}{|\lambda|}\right), \quad (2.2)$$

$$\vartheta'(x, \lambda, q) = -\sqrt{\lambda} \sin \sqrt{\lambda}x + \frac{1}{2} \int_0^x \left(\cos \sqrt{\lambda}x + \cos \sqrt{\lambda}(x-2t) \right) q(t)dt + O\left(\frac{e^{|\operatorname{Im} \sqrt{\lambda}|x}}{|\lambda|^{1/2}}\right), \quad (2.3)$$

$$\varphi(x, \lambda, q) = \frac{\sin \sqrt{\lambda}x}{\sqrt{\lambda}} + \frac{1}{2\lambda} \int_0^x \left(-\cos \sqrt{\lambda}x + \cos \sqrt{\lambda}(x-2t) \right) q(t)dt + O\left(\frac{e^{|\operatorname{Im} \sqrt{\lambda}|x}}{|\lambda|^{3/2}}\right), \quad (2.4)$$

$$\varphi'(x, \lambda, q) = \cos \sqrt{\lambda}x + \frac{1}{2\sqrt{\lambda}} \int_0^x \left(\sin \sqrt{\lambda}x - \sin \sqrt{\lambda}(x-2t) \right) q(t)dt + O\left(\frac{e^{|\operatorname{Im} \sqrt{\lambda}|x}}{|\lambda|}\right) \quad (2.5)$$

as $|\lambda| \rightarrow \infty$, uniformly on bounded sets of $(x; q) \in [0, 1] \times L^2_C(0, 1)$.

Lemma 2.1. *For each $(q; b) \in L^2(0, 1) \times \mathbb{R}$ and $n \geq 0$ the following identities are fulfilled:*

$$\|\varphi(\cdot, \lambda_n, q)\|^2 = (-1)^{n+1} e^{\nu_n} \dot{w}(\lambda_n, q, b), \quad \|\xi_b(\cdot, \lambda_n, q)\|^2 = (-1)^{n+1} e^{-\nu_n} \dot{w}(\lambda_n, q, b),$$

$$\psi_n^2(x, q, b) = \frac{\varphi(x, \lambda_n, q) \xi_b(x, \lambda_n, q)}{\dot{w}(\lambda_n, q, b)},$$

where $\lambda_n = \lambda_n(q, b)$ and $\nu_n = \nu_n(q, b)$.

Proof. Let $\varphi(x) = \varphi(x, \lambda_n, q)$, $\xi_b(x) = \xi_b(x, \lambda_n, q)$ and so on. Definition (1.3) yields

$$\varphi(x) = -\varphi(1) \xi_b(x) = (-1)^{n+1} e^{\nu_n} \xi_b(x).$$

This identity and $\varphi^2 = \{\dot{\varphi}, \varphi\}'$ give

$$\|\varphi\|^2 = \{\dot{\varphi}, \varphi\}(1) = (-1)^{n+1} e^{\nu_n} (\dot{\varphi} \xi'_b - \dot{\varphi}' \xi_b)(1) = (-1)^{n+1} e^{\nu_n} \dot{w}(\lambda_n, q, b),$$

since $\dot{\varphi}(0) = \dot{\varphi}'(0) = \dot{\xi}_b(1) = \dot{\xi}'_b(1) = 0$. Furthermore, we have

$$\psi_n^2(x) = \frac{\varphi^2(x)}{\|\varphi\|^2} = \frac{\varphi(x) \cdot (-1)^{n+1} e^{-\nu_n} \varphi(x)}{\dot{w}(\lambda_n, q, b)} = \frac{\varphi(x) \xi_b(x)}{\dot{w}(\lambda_n, q, b)}$$

and $\|\xi_b\|^2 = e^{-2\nu_n} \|\varphi\|^2 = (-1)^{n+1} e^{-\nu_n} \dot{w}(\lambda_n, q, b)$. □

Lemma 2.2. For each $(q; b) \in L^2(0, 1) \times \mathbb{R}$ the identity

$$b = \sum_{n \geq 0} \left(2 - \frac{e^{\nu_n}}{|\dot{w}(\lambda_n, q, b)|} \right)$$

holds true, where $\lambda_n = \lambda_n(q, b)$ and $\nu_n = \nu_n(q, b)$, $n \geq 0$.

Proof. Consider the meromorphic function

$$f(\lambda) = \frac{\varphi(1, \lambda, q)}{w(\lambda, q, b)} = \frac{\varphi}{\varphi' + b\varphi}(1, \lambda, q), \quad \lambda \in \mathbb{C}.$$

All roots $\lambda_n = \lambda_n(q, b)$ of $w(\lambda) = w(\lambda, q, b)$ are simple and $\text{sign } \dot{w}(\lambda_n) = (-1)^{n+1}$. Therefore, definition (1.3) gives

$$\text{res}_{\lambda=\lambda_n} f(\lambda) = \frac{\varphi(1, \lambda, q)}{\dot{w}(\lambda_n)} = \frac{(-1)^n e^{\nu_n}}{\dot{w}(\lambda_n)} = -\frac{e^{\nu_n}}{|\dot{w}(\lambda_n)|}, \quad n \geq 0, \quad (2.6)$$

Put $|\lambda| = \pi^2 m^2 \rightarrow \infty$. Then, due to (2.4), (2.5) and $\int_0^1 e^{ik(1-2t)} q(t) dt = o(e^{|\text{Im } k|})$, $|k| \rightarrow \infty$, we have

$$\begin{aligned} f(\lambda) &= \frac{\lambda^{-1/2} \sin \sqrt{\lambda} - \lambda^{-1} \cos \sqrt{\lambda} \cdot \frac{1}{2} Q_0 + o(\lambda^{-1} e^{|\text{Im } \sqrt{\lambda}|})}{\cos \sqrt{\lambda} + \lambda^{-1/2} \sin \sqrt{\lambda} \cdot (\frac{1}{2} Q_0 + b) + o(\lambda^{-1/2} e^{|\text{Im } \sqrt{\lambda}|})} \\ &= \frac{\lambda^{-1/2} \tan \sqrt{\lambda} - \lambda^{-1} \cdot \frac{1}{2} Q_0 + o(\lambda^{-1})}{1 + \lambda^{-1/2} \tan \sqrt{\lambda} \cdot (\frac{1}{2} Q_0 + b) + o(\lambda^{-1/2})} \\ &= \lambda^{-\frac{1}{2}} \tan \sqrt{\lambda} - \lambda^{-1} \cdot \frac{1}{2} Q_0 - \lambda^{-1} \tan^2 \sqrt{\lambda} \cdot (\frac{1}{2} Q_0 + b) + o(\lambda^{-1}). \end{aligned}$$

Let $f_0(\lambda) = \lambda^{-\frac{1}{2}} \tan \sqrt{\lambda}$. Applying the Cauchy Theorem in the disk $|\lambda| < \pi^2 m^2$, we obtain

$$\sum_{n=0}^m \left(\text{res}_{\lambda=\lambda_n} f(\lambda) - \text{res}_{\lambda=\lambda_n^0} f_0(\lambda) \right) = -\frac{1}{2} Q_0 - (\frac{1}{2} Q_0 + b) \cdot \sum_{n=0}^m \text{res}_{\lambda=\lambda_n^0} (f_0(\lambda))^2. \quad (2.7)$$

Note that

$$f_0(\lambda) = \frac{\tan \sqrt{\lambda}}{\sqrt{\lambda}} = -\frac{2}{\lambda - \lambda_n^0} \left(1 + \frac{\lambda - \lambda_n^0}{4\lambda_n^0} + O((\lambda - \lambda_n^0)^2) \right) \quad \text{as } \lambda \rightarrow \lambda_n^0, \quad n \geq 0.$$

Hence,

$$\text{res}_{\lambda=\lambda_n^0} f_0(\lambda) = -2, \quad \text{res}_{\lambda=\lambda_n^0} (f_0(\lambda))^2 = -\frac{2}{\lambda_n^0}, \quad n \geq 0,$$

and

$$\sum_{n \geq 0} \text{res}_{\lambda=\lambda_n^0} (f_0(\lambda))^2 = -\frac{2}{\pi^2} \sum_{n \geq 0} \frac{1}{(n + \frac{1}{2})^2} = -\frac{2}{\pi^2} \cdot \frac{\pi^2}{2} = -1.$$

Substituting these identities and (2.6) into (2.7), we obtain (1.4) as $m \rightarrow \infty$. $\square$

Theorem 2.3 (Uniqueness Theorem). *Let $\lambda_n(q, b) = \lambda_n(p, h)$ and $\nu_n(q, b) = \nu_n(p, h)$ for all $n \geq 0$ and some $(q; b), (p; h) \in L^2(0, 1) \times \mathbb{R}$. Then $(q; b) = (p; h)$.*

Proof. Lemma 2.2 immediately gives $b = h$. The rest of the proof is standard (see p. 58 [PT]). Define functions

$$f_1(\lambda, x) = \frac{F_1(\lambda, x)}{w(\lambda, q, b)}, \quad F_1(\lambda; x) = \varphi(x, \lambda, p)\xi'_b(x, \lambda, q) - \xi_b(x, \lambda, p)\varphi'(x, \lambda, q),$$

$$f_2(\lambda, x) = \frac{F_2(\lambda, x)}{w(\lambda, q, b)}, \quad F_2(\lambda, x) = \varphi(x, \lambda, p)\xi_b(x, \lambda, q) - \xi_b(x, \lambda, p)\varphi(x, \lambda, q).$$

Recall that

$$w(\lambda, q, b) = \varphi'(1, \lambda, q) + b\varphi(1, \lambda, q) = \{\varphi, \xi_b\}(\lambda, q).$$

All roots $\lambda_n(q, b) = \lambda_n(p, b)$ of the denominator $w(\cdot, q, b)$ are simple. Moreover, all these values are roots of the numerators F_1, F_2 , since definition (1.3) of the norming constant ν_n yields

$$\varphi(x, \lambda_n, p) = (-1)^{n+1}e^{\nu_n}\xi_b(x, \lambda_n, p), \quad \varphi(x, \lambda_n, q) = (-1)^{n+1}e^{\nu_n}\xi_b(x, \lambda_n, q).$$

Hence, the functions f_1, f_2 are entire with respect to λ for each $x \in [0, 1]$. Let $|\lambda| = \pi^2 m^2$, $m \rightarrow \infty$. Due to asymptotics (2.4), (2.5), we have

$$w(\lambda, q, b) = \cos \sqrt{\lambda} + O(|\lambda|^{-1/2}e^{|\operatorname{Im} \sqrt{\lambda}|}) = \cos \sqrt{\lambda} \cdot (1 + O(|\lambda|^{-1/2})).$$

Note that

$$\xi_b(x, \lambda, q) = -\vartheta(1 - x, \lambda, q^*) - b\varphi(1 - x, \lambda, q^*),$$

where $q^*(x) = q(1 - x)$, $x \in [0, 1]$. Therefore, asymptotics (2.2)-(2.5) give

$$F_1(x, \lambda) = -\sin \sqrt{\lambda}x \cdot \sin \sqrt{\lambda}(1 - x) + \cos \sqrt{\lambda}x \cdot \cos \sqrt{\lambda}(1 - x) + O(|\lambda|^{-1/2}e^{|\operatorname{Im} \sqrt{\lambda}|})$$

$$= \cos \sqrt{\lambda} \cdot (1 + O(|\lambda|^{-1/2})) \quad \text{and} \quad F_2(x, \lambda) = O(|\lambda|^{-1/2}e^{|\operatorname{Im} \sqrt{\lambda}|}).$$

Thus,

$$f_1(x, \lambda) = 1 + O(|\lambda|^{-1/2}) \quad \text{and} \quad f_2(x, \lambda) = O(|\lambda|^{-1/2}) \quad \text{as} \quad |\lambda| = \pi^2 m^2 \rightarrow \infty.$$

Since f_1 and f_2 are entire with respect to λ , the maximum principle yields $f_1(x, \lambda) = 1$ and $f_2(x, \lambda) = 0$ for each $x \in [0, 1]$ and $\lambda \in \mathbb{C}$. In other words,

$$(\varphi(x, \lambda, p) - \varphi(x, \lambda, q))\xi'_b(x, \lambda, q) - (\xi_b(x, \lambda, p) - \xi_b(x, \lambda, q))\varphi'(x, \lambda, q) = 0,$$

$$(\varphi(x, \lambda, p) - \varphi(x, \lambda, q))\xi_b(x, \lambda, q) - (\xi_b(x, \lambda, p) - \xi_b(x, \lambda, q))\varphi(x, \lambda, q) = 0.$$

This gives $\varphi(x, \lambda, p) = \varphi(x, \lambda, q)$ for all $x \in [0, 1]$ and $\lambda \neq \lambda_n(q)$, $n \geq 0$. Hence, $p = q$. $\square$

3 Analyticity, asymptotics and local isomorphism

Lemma 3.1. *Let $(q; b) \in L_{\mathbb{C}}^2(0, 1) \times \mathbb{C}$ and $N > 2(\|q\| + |b|)e^{\|q\|}$. Then $w(\lambda)$ has exactly N roots, counted with multiplicities, in the disc $\{\lambda : |\lambda| < \pi^2 N^2\}$ and for each $n > N$ exactly one simple root $\lambda_n(q, b)$ in the region $\{\lambda : |\sqrt{\lambda} - k_n^0| < \pi/4\}$. There are no other roots. Moreover, the following asymptotics is fulfilled:*

$$\lambda_n(q, b) = \lambda_n^0 + O(1) \quad \text{as } n \rightarrow \infty, \quad (3.1)$$

uniformly on bounded subsets of $L_{\mathbb{C}}^2(0, 1) \times \mathbb{C}$.

Proof. The standard estimates of φ and φ' (see p. 13 [PT]) give

$$|\varphi'(1, \lambda, q) - \cos \sqrt{\lambda}| \leq |\lambda|^{-\frac{1}{2}} \cdot \|q\| e^{\|q\| + |\operatorname{Im} \sqrt{\lambda}|}, \quad |\varphi(1, \lambda, q)| \leq |\lambda|^{-\frac{1}{2}} \cdot e^{\|q\| + |\operatorname{Im} \sqrt{\lambda}|}.$$

Hence,

$$|w(\lambda, q, b) - \cos \sqrt{\lambda}| \leq |\lambda|^{-\frac{1}{2}} \cdot (\|q\| + |b|) e^{\|q\| + |\operatorname{Im} \sqrt{\lambda}|} < \frac{1}{2} N |\lambda|^{-1/2} e^{|\operatorname{Im} \sqrt{\lambda}|}.$$

Let $\lambda \in L_N \cup \bigcup_{n>N} l_n$, where $L_N = \{\lambda : |\lambda| = \pi^2 N^2\}$ and $l_n = \{\lambda : |\sqrt{\lambda} - k_n^0| = \pi/4\}$, $n > N$. Then the simple estimate $4|\cos \sqrt{\lambda}| > e^{|\operatorname{Im} \sqrt{\lambda}|}$ is fulfilled (p. 27 [PT]) and we have

$$|w(\lambda, q, b) - \cos \sqrt{\lambda}| < 2N |\lambda|^{-1/2} \cdot |\cos \sqrt{\lambda}| < |\cos \sqrt{\lambda}|, \quad \lambda \in L_N \cup \bigcup_{n>N} l_n.$$

Therefore, by Rouché's Theorem, $w(\cdot, q, b)$ has as many roots as $\cos \sqrt{\lambda}$ in each region bounded by these contours and the remaining unbounded domain. Furthermore, it follows from (2.4), (2.5) that $0 = w(\lambda_n, q, b) = \cos \sqrt{\lambda_n} + O(n^{-1})$ as $n \rightarrow \infty$, uniformly on bounded subsets of $L_{\mathbb{C}}^2(0, 1) \times \mathbb{C}$. Note that the point k_n^0 is the simple root of the function $\cos z$. Hence, we have $\sqrt{\lambda_n} = k_n^0 + O(n^{-1})$ or, equivalently, (3.1). $\square$

Define the inner product in $L_{\mathbb{C}}^2(0, 1) \times \mathbb{C}$ by

$$\langle (q; b), (p; h) \rangle = \int_0^1 q(t) \overline{p(t)} dt + b \bar{h}.$$

Below we write $d_{(q;b)}\sigma(q, b) = (f; g)$ iff the identity $[d_{(q;b)}\sigma(q, b)](p; h) = \langle (f; g), (p; h) \rangle$ is fulfilled for all $(p; h) \in L^2(0, 1) \times \mathbb{R}$. We use the similar notation $\partial_{(q;b)}\sigma(\lambda, q, b)$ for the partial derivative of σ with respect to $(q; b)$.

Lemma 3.2. *(i) All functions $\lambda_n(q, b)$, $\mu_n(q, b)$, $\nu_n(q, b)$, $n \geq 0$, are real-analytic on $L^2(0, 1) \times \mathbb{R}$. Their derivatives are given by*

$$d_{(q(x);b)}\lambda_n(q, b) = (\psi_n^2(x); \psi_n^2(1)), \quad d_{(q(x);b)}\mu_n(q, b) = (\psi_n^2(x) - 1; \psi_n^2(1) - 2), \quad (3.2)$$

$$d_{(q(x);b)}\nu_n(q, b) = ((\psi_n \chi_n)(x); (\psi_n \chi_n)(1)), \quad (3.3)$$

where $\psi_n(x) = \psi_n(x, q, b)$ is the n -th normalized eigenfunction of the operator H and

$$\chi_n(x) = \chi_n(x, q, b) = \frac{\vartheta(x, \lambda_n, q)}{\psi_n'(0, q, b)} - \psi_n(x, q, b) \int_0^1 (\varphi \vartheta)(t, \lambda_n, q) dt. \quad (3.4)$$

(ii) Let $(q; b) \in L^2(0, 1) \times \mathbb{R}$. Then, each function λ_n, μ_n, ν_n , has an analytic continuation from $L^2(0, 1) \times \mathbb{R}$ into some complex ball $\{(p; h) \in L^2_{\mathbb{C}}(0, 1) \times \mathbb{C} : \|(p - q; b - h)\| < \varepsilon\}$, where $\varepsilon = \varepsilon(q, b) > 0$ doesn't depend on n .

Remark 3.3. Since $\psi_n(0) = 0$, we have $\psi_n(x) = \psi'_n(0)\varphi(x)$. Therefore, $\chi_n(x)$ is the solution of (2.1) for $\lambda = \lambda_n$ such that $\{\chi_n, \psi_n\} = 1$ and $\int_0^1 \psi_n(x)\chi_n(x)dx = 0$. Note that these conditions define χ_n uniquely.

Proof of Lemma 3.2. (i) It is well-known (see p. 21 [PT]) that

$$\begin{aligned}\partial_{q(x)}\varphi(1, \lambda, q) &= \varphi(x)(\vartheta(x)\varphi(1) - \varphi(x)\vartheta(1)), \\ \partial_{q(x)}\varphi'(1, \lambda, q) &= \varphi(x)(\vartheta(x)\varphi'(1) - \varphi(x)\vartheta'(1)),\end{aligned}$$

where $\varphi(x) = \varphi(x, \lambda, q)$, $\vartheta(x) = \vartheta(x, \lambda, q)$ and so on. This gives

$$\partial_{(q(x); b)}w(\lambda, q, b) = ((\varphi' + b\varphi)(1)\vartheta(x) - (\vartheta' + b\vartheta)(1)\varphi(x))\varphi(x); \varphi(1)).$$

Note that

$$(\varphi' + b\varphi)(1)\vartheta(x) - (\vartheta' + b\vartheta)(1)\varphi(x) = -\xi_b(x),$$

since both functions are solutions of (2.1) with the same initial data at $x = 1$. Therefore,

$$\partial_{(q(x); b)}w(\lambda, q, b) = (-\varphi(x)\xi_b(x); -\varphi(1)\xi_b(1)).$$

Recall that the function $w(\cdot, \cdot, \cdot)$ is entire and $w(\lambda_n(q, b), q, b) = 0$. Then, the Implicit Function Theorem and Lemma 2.1 give

$$d_{(q(x); b)}\lambda_n(q, b) = -\frac{\partial_{(q(x); b)}w(\lambda_n, q, b)}{w(\lambda_n, q, b)} = (\psi_n^2(x); \psi_n^2(1)).$$

The identity $\mu_n(q, b) = \lambda_n(q, b) - \lambda_n^0 - \int_0^1 q(t)dt - 2b$ yields

$$d_{(q(x); b)}\mu_n(q, b) = d_{(q(x); b)}\lambda_n(q, b) - (1; 2) = (\psi_n^2(x) - 1; \psi_n^2(1) - 2).$$

Furthermore, definition (1.3) implies

$$d_{(q(x); b)}\nu_n(q, b) = \frac{d_{(q(x); b)}\varphi(1, \lambda_n(q, b), q)}{\varphi(1)} = \frac{(\partial_{q(x)}\varphi(1, \lambda_n, q); 0) + \dot{\varphi}(1)d_{(q(x); b)}\lambda_n(q, b)}{\varphi(1)}.$$

Substituting $\partial_{q(x)}\varphi(1, \lambda_n, q)$ and $d_{(q(x); b)}\lambda_n(q, b)$ into this formula and using the identity $\psi_n(x) = \psi'_n(0)\varphi(x)$, we obtain

$$d_{(q(x); b)}\nu_n(q, b) = (\psi_n(x)\chi_n(x); \psi_n(1)\chi_n(1)), \quad \chi_n(x) = \frac{\vartheta(x, \lambda_n, q)}{\psi'_n(0, q, b)} - C_n\psi_n(x, q, b),$$

where $C_n = C_n(q, b)$ is some constant. In order to find C_n , note that $\nu_n(q+c, b) = \nu_n(q, b)$ for all $c \in \mathbb{R}$. Therefore,

$$0 = \langle \partial_{q(x)}\nu_n(q, b), 1 \rangle = \int_0^1 \psi_n(x)\chi_n(x)dx = \int_0^1 \varphi(x)\vartheta(x)dx - C_n \int_0^1 \psi_n^2(x)dx.$$

Since $\int_0^1 \psi_n^2(x)dx = 1$, this yields $C_n = \int_0^1 \varphi(x)\vartheta(x)dx$.

(ii) The proof repeats p. 51 [PT] for eigenvalues and p. 64 for norming constants. $\square$

Lemma 3.4. (i) *The following asymptotics are fulfilled:*

$$\psi_n(x, q, b) = \sqrt{2} \sin k_n^0 x + O(n^{-1}), \quad \chi_n(x, q, b) = \frac{\cos k_n^0 x}{\sqrt{2} k_n^0} + O(n^{-2}), \quad n \rightarrow \infty, \quad (3.5)$$

$$\mu_n(q, b) = -\hat{q}_{n+\frac{1}{2}}^{(cos)} + O(n^{-1}), \quad \nu_n(q, b) = \nu_n^0 + \frac{\hat{q}_{n+\frac{1}{2}}^{(sin)}}{2k_n^0} + O(n^{-2}), \quad n \rightarrow \infty, \quad (3.6)$$

uniformly on bounded subsets of $[0, 1] \times L_{\mathbb{C}}^2(0, 1) \times \mathbb{C}$, where

$$\hat{q}_{n+\frac{1}{2}}^{(cos)} = \int_0^1 q(x) \cos 2k_n^0 x dx, \quad \hat{q}_{n+\frac{1}{2}}^{(sin)} = \int_0^1 q(x) \sin 2k_n^0 x dx.$$

(ii) *The mapping Φ is real-analytic on $L^2(0, 1) \times \mathbb{R}$ and its Fréchet derivative is given by*

$$[d_{(q;b)}\Phi](\cdot) = \left(\langle (1; 2), \cdot \rangle; \{ \langle X_n, \cdot \rangle \}_0^\infty; \{ \langle Y_n, \cdot \rangle \}_0^\infty \right), \quad (3.7)$$

$$X_n = (\psi_n^2(x) - 1; \psi_n^2(1) - 2), \quad Y_n = ((\psi_n \chi_n)(x); (\psi_n \chi_n)(1)), \quad n \geq 0.$$

Proof. (i) Let $\lambda_n = \lambda_n(q, b)$. Due to (3.1), we have $\sqrt{\lambda_n} = k_n^0 + O(n^{-1})$. Asymptotics (2.2)-(2.5) yields

$$\varphi(x, \lambda_n, q) = \frac{\sin k_n^0 x}{k_n^0} + O(n^{-2}), \quad \|\varphi(\cdot, \lambda_n, q)\|^2 = \int_0^1 \varphi^2(x, \lambda_n, q) dx = \frac{1}{2\lambda_n^0} + O(n^{-3}).$$

Hence,

$$\psi_n(x, q, b) = \frac{\varphi(x, \lambda_n, q)}{\|\varphi(\cdot, \lambda_n, q)\|} = \sqrt{2} \sin k_n^0 x + O(n^{-1}).$$

This gives $\psi'_n(0, q, b) = \|\varphi(\cdot, \lambda_n, q)\|^{-1} = \sqrt{2} k_n^0 + O(1)$. Also,

$$\vartheta(x, \lambda_n, q) = \cos k_n^0 x + O(n^{-1}), \quad \int_0^1 (\vartheta \varphi)(x, \lambda_n, q) dx = O(n^{-2}).$$

Using definition (3.4), we obtain the second asymptotics in (3.5). Lemma 3.2 and (3.5) yield

$$\nu_n(q, b) = \int_0^1 [d_{(q;b)}\mu_n(tq, tb)](q; b) dt = \lambda_n^0 + \langle (2 \sin^2 k_n^0 x - 1; 0), (q(x); b) \rangle + O(n^{-1}),$$

$$\nu_n(q, b) = \nu_n^0 + \int_0^1 [d_{(q;b)}\nu_n(tq, tb)](q; b) dt = \nu_n^0 + \frac{\langle (\sin k_n^0 x \cos k_n^0 x; 0), (q(x); b) \rangle}{k_n^0} + O(n^{-2}).$$

This gives (3.6) since $2 \sin^2 k_n^0 x = 1 - \cos 2k_n^0 x$ and $\sin k_n^0 x \cos k_n^0 x = \frac{1}{2} \sin 2k_n^0 x$.

(ii) As it shown above, $\Phi : L^2(0, 1) \times \mathbb{R} \rightarrow \mathbb{R} \times \mathcal{M} \times \ell_1^2$ is locally bounded and all "coordinate functions" $\int_0^1 q(t) dt + 2b$, $\lambda_n(q, b)$, $\nu_n(q, b) - \nu_n^0$ are real-analytic. Therefore (e.g., see p. 138 [PT]), Φ is a real-analytic mapping and its Fréchet derivative is given by (3.7). $\square$

Lemma 3.5. *Let $(q; b) \in L^2(0, 1) \times \mathbb{R}$ and*

$$X_n = (\psi_n^2(x) - 1; \psi_n^2(1) - 2), \quad Y_n = ((\psi_n \chi_n)(x); (\psi_n \chi_n)(1)), \quad n \geq 0,$$

$$Z_m = (-2(\psi_m \chi_m)'(x); (\psi_m \chi_m)(1)), \quad T_m = (2(\psi_m^2)'(x); -\psi_m^2(1)), \quad m \geq 0.$$

Then vectors $(1; 0)$, $\{Z_m\}_0^\infty$, $\{T_m\}_0^\infty$ form the biorthogonal basis to $(1; 2)$, $\{X_n\}_0^\infty$, $\{Y_n\}_0^\infty$. In other words,

$$\begin{aligned} \langle (1; 2), (1; 0) \rangle &= 1, & \langle (1; 2), Z_m \rangle &= 0, & \langle (1; 2), T_m \rangle &= 0 \\ \langle X_n, (1; 0) \rangle &= 0, & \langle X_n, Z_m \rangle &= \delta_{nm}, & \langle X_n, T_m \rangle &= 0, \\ \langle Y_n, (1; 0) \rangle &= 0, & \langle Y_n, Z_m \rangle &= 0, & \langle Y_n, T_m \rangle &= \delta_{nm}, \end{aligned} \quad n, m \geq 0.$$

Proof. By definition, $\langle (1; 2), (1; 0) \rangle = 1$ and $\langle (1; 2), Z_m \rangle = \langle (1; 2), T_m \rangle = 0$, since $\psi_m(0) = 0$. Due to $\int_0^1 \psi_n^2(x) dx = 1$ and $\int_0^1 (\psi_n \chi_n)(x) dx = 0$ (see Remark 3.3), we obtain $\langle X_n, (1; 0) \rangle = \langle Y_n, (1; 0) \rangle = 0$. We consider $\langle X_n, Z_m \rangle$. The partial integration gives

$$\begin{aligned} \langle X_n, Z_m \rangle &= -2 \int_0^1 (\psi_n^2(x) - 1)(\psi_m \chi_m)'(x) dx + (\psi_n^2(1) - 2)(\psi_m \chi_m)(1) \\ &= -2 \int_0^1 \psi_n^2(x)(\psi_m \chi_m)'(x) dx + (\psi_n^2(1) - 2)(\psi_m \chi_m)(1) = - \int_0^1 (\psi_n^2(\psi_m \chi_m)' - (\psi_n^2)' \psi_m \chi_m)(x) dx \\ &= - \int_0^1 (\{\psi_n, \psi_m\} \cdot \psi_n \chi_m + \psi_n \psi_m \cdot \{\psi_n, \chi_m\})(x) dx. \end{aligned}$$

If $n \neq m$, then we have $\{\psi_n, \psi_m\}' = (\lambda_n - \lambda_m)\psi_n \psi_m$, $\{\psi_n, \chi_m\}' = (\lambda_n - \lambda_m)\psi_n \chi_m$. Hence,

$$\langle X_n, Z_m \rangle = - \frac{\{\psi_n, \psi_m\} \{\psi_n, \chi_m\}}{\lambda_n - \lambda_m} \Big|_0^1 = 0, \quad n \neq m,$$

since $\{\psi_n, \psi_m\}(0) = \{\psi_n, \psi_m\}(1) = 0$. If $n = m$, then $\{\psi_n, \psi_m\} = 0$, $\{\psi_n, \chi_n\} = -1$ and

$$\langle X_n, Z_n \rangle = \int_0^1 \psi_n^2(x) dx = 1, \quad n \geq 0.$$

The proof of other identities is similar. □

Theorem 3.6 (Local Isomorphism). *For each $(q; b) \in L^2(0, 1) \times \mathbb{R}$ operator $d_{(q; b)} \Phi : L^2(0, 1) \times \mathbb{R} \rightarrow \mathbb{R} \times \ell^2 \times \ell_1^2$ given by (3.7) is invertible.*

Proof. Due to Lemma 3.4, we have

$$X_n = (-\cos \pi(2n+1)x; 0) + O(n^{-1}), \quad 2k_n^0 Y_n = (\sin \pi(2n+1)x; 0) + O(n^{-1}), \quad n \geq 0.$$

Note that vectors $(0; 1)$, $\{(2 \cos \pi(2n+1)x; 0)\}_0^\infty$, $\{(2 \sin \pi(2n+1)x; 0)\}_0^\infty$ form the orthonormal basis in $L^2(0, 1) \times \mathbb{R}$ and the error terms are square summable. Then, $d_{(q; b)} \Phi$ is a Fredholm operator. Due to Lemma 3.5, the vectors $(1; 2)$, $\{X_n\}_0^\infty$, $\{Y_n\}_0^\infty$ are linearly independent. Using the standard arguments from the functional analysis (e.g., see p.163 [PT]), we deduce that $(d_{(q; b)} \Phi)^{-1}$ is bounded. □

4 Proof of Theorem 1.1 and Corollaries 1.3, 1.2

Lemma 4.1. *For each $(c^*, \{\mu_n^*\}_0^\infty) \in \mathbb{R} \times \mathcal{M}$ there exists a potential $q^* \in L^2(0, 1)$ such that $\lambda_n(q^*, 0) = \lambda_n^0 + c^* + \mu_n^*$ for all $n \geq 0$.*

Proof. Define $\lambda_{n+\frac{1}{2}}^0$ and $\lambda_{n+\frac{1}{2}}^*$ by

$$\lambda_{n+\frac{1}{2}}^0 = \pi^2(n+1)^2, \quad \lambda_{n+\frac{1}{2}}^* = \frac{\lambda_{n+1}^0 - \lambda_{n+\frac{1}{2}}^0}{\lambda_{n+1}^0 - \lambda_n^0} \cdot \lambda_n^* + \frac{\lambda_{n+\frac{1}{2}}^0 - \lambda_n^0}{\lambda_{n+1}^0 - \lambda_n^0} \cdot \lambda_{n+1}^*, \quad n \geq 0.$$

Then, $\lambda_n^* < \lambda_{n+\frac{1}{2}}^* < \lambda_{n+1}^*$, $n \geq 0$, and $\lambda_{m/2}^* = \pi^2(m/2)^2 + c^* + \mu_{m/2}^*$, where $\{\mu_{m/2}^*\}_0^\infty \in \ell^2$. Using [PT], we see that there exists an "even" potential $q \in L^2(0, 2)$, $q(x) = q(2-x)$, $x \in [0, 2]$, such that $\{\lambda_{m/2}^*\}_0^\infty$ is the Dirichlet spectrum of q on the interval $[0, 2]$. Put $q^* = q|_{[0,1]}$. Then we have $\lambda_n(q^*, 0) = \lambda_{2n/2}^* = \lambda_n^*$ for all $n \geq 0$. $\square$

Lemma 4.2. *Fix any $(q; b) \in L^2(0, 1) \times \mathbb{R}$, $n \geq 0$ and $t \in \mathbb{R}$. Denote*

$$q_n^t(x) = q(x) - 2 \frac{d^2}{dx^2} \log \eta_n^t(x, q, b), \quad \text{where } \eta_n^t(x, q, b) = 1 + (e^t - 1) \int_x^1 \psi_n^2(t, q, b) dt, \quad x \in [0, 1],$$

$$b_n^t = b - (e^t - 1) \psi_n^2(1, q, b).$$

Then $(q_n^t; b_n^t) \in L^2(0, 1) \times \mathbb{R}$ and

$$\lambda_m(q_n^t, b_n^t) = \lambda_m(q, b), \quad \nu_m(q_n^t, b_n^t) = \nu_m(q, b) + t \delta_{mn} \quad \text{for all } m \geq 0.$$

Proof. Let $\psi_n(x) = \psi_n(x, q, b)$ and so on. Repeating the arguments of p. 91–93 [PT] or using direct calculations, it is easy to check that for each $m \geq 0$ the function

$$\tilde{\psi}_m(x) = \psi_m(x) - (e^t - 1) \frac{\psi_n(x)}{\eta_n^t(x)} \int_x^1 \psi_n(t) \psi_m(t) dt,$$

is some solutions of $-\psi'' + q_n^t(x)\psi = \lambda_m(q, b)\psi$ (in particular, $\tilde{\psi}_n(x) = \psi_n(x)/\eta_n^t(x)$). Since $\psi_m(0) = \psi_n(0) = 0$, we have $\tilde{\psi}_m(0) = 0$, $m \geq 0$. The crucial point is the new boundary condition at $x = 1$. We have

$$\tilde{\psi}_m'(1) + b_n^t \tilde{\psi}_m(1) = \psi_m'(1) + (e^t - 1) \psi_n^2(1) \psi_m(1) + b_n^t \tilde{\psi}_m(1) = \psi_m'(1) + b \psi_m(1) = 0$$

for all $m \geq 0$. Therefore, $\lambda_m(q_n^t, b_n^t) = \lambda_m(q, b)$, $m \geq 0$ (there are no other roots due to Lemma 3.1). Furthermore,

$$\tilde{\psi}_m'(0) = \psi_m'(0), \quad m \neq n, \quad \text{and} \quad \tilde{\psi}_n'(0) = e^{-t} \psi_n'(0).$$

By definition (1.3), it gives $\nu_m(q_n^t, b_n^t) = \nu_m(q, b) + t \delta_{mn}$ for all $m \geq 0$. $\square$

Proof of Theorem 1.1. ii) Identity (1.4) was proved in Lemma 2.2.

(i) It follows from Lemma 3.4 that Φ maps $L^2(0, 1) \times \mathbb{R}$ into $\mathbb{R} \times \mathcal{M} \times \ell_1^2$ and Φ is real-analytic and its Fréchet derivative is given by (3.7). Theorem 3.6 yields that Φ is a local real-analytic isomorphism. Furthermore, due to Theorem 2.3, Φ is one-to-one. We prove that Φ is onto.

Let $c^* \in \mathbb{R}$, $\{\mu_n^*\}_0^\infty \in \mathcal{M}$ and $\{\nu_n^* - \nu_n^0\}_0^\infty \in \ell_1^2$ be an arbitrary spectral data. Due to Lemma 4.1, there exists a potential $q^* \in L^2(0, 1)$ such that $\lambda_n(q^*, 0) = \lambda_n^* = \lambda_n^0 + c^* + \mu_n^*$ for all $n \geq 0$. Let $\eta_n = \nu_n(q^*, 0) - \nu_n^0$. Note that we have the following convergence:

$$(\eta_0, \dots, \eta_{N-1}, \nu_N^* - \nu_N^0, \nu_{N+1}^* - \nu_{N+1}^0, \dots) \rightarrow (\eta_0, \dots, \eta_{N-1}, \eta_N, \dots) \text{ in } \ell_1^2 \text{ as } N \rightarrow \infty.$$

Theorem 3.6 yields that the mapping Φ is invertible in some neighborhood of the point $\Phi(q^*, 0) = (c^*; \{\mu_n^*\}_0^\infty; \{\eta_n\}_0^\infty) \in \mathbb{R} \times \mathcal{M} \times \ell_1^2$. Hence, for some large integer N and $(q^{(N)}; b^{(N)}) \in L^2(0, 1) \times \mathbb{R}$ we have

$$(c^*; \{\mu_n^*\}_0^\infty; (\eta_0, \dots, \eta_{N-1}, \nu_N^* - \nu_N^0, \nu_{N+1}^* - \nu_{N+1}^0, \dots)) = \Phi(q^{(N)}, b^{(N)}).$$

Applying Lemma 4.2 step by step, we construct the sequence of potentials and boundary constants

$$(q^{(k)}; b^{(k)}) = (q^{(k+1)}; b^{(k+1)})_{t_k}^{t_k} \in L^2(0, 1) \times \mathbb{R}, \quad t_k = \nu_k^* - \nu_k^0 - \eta_k, \quad k = N-1, N-2, \dots, 0,$$

such that $\Phi(q^{(k)}, b^{(k)}) = (c^*; \{\mu_n^*\}_0^\infty; (\eta_0, \dots, \eta_{k-1}, \nu_k^* - \nu_k^0, \nu_{k+1}^* - \nu_{k+1}^0, \dots))$. In particular, $\lambda_n(q^{(0)}, b^{(0)}) = \lambda_n^*$ and $\nu_n(q^{(0)}, b^{(0)}) = \nu_n^*$ for all $n \geq 0$. We are done. $\square$

Proof of Corollary 1.3. Fix some $b, c^* \in \mathbb{R}$, $\{\mu_n^*\}_0^\infty \in \mathcal{M}$ and $m \geq 0$. It follows from Theorem 1.1 that the mapping $q \mapsto \{\nu_n(q, b) - \nu_n^0\}_0^\infty$ is a real-analytic isomorphism between $\text{Iso}_b(\{\lambda_n^*\}_0^\infty)$ and the set

$$\mathcal{N}^b = \left\{ \{\nu_n - \nu_n^0\}_0^\infty \in \ell_1^2 : \sum_{n \geq 0} \left(2 - \frac{e^{\nu_n}}{|\dot{w}^*(\lambda_n^*)|} \right) = b \right\} \subset \ell_1^2.$$

Introduce the mapping $P_m : \{\nu_n - \nu_n^0\}_0^\infty \mapsto \{\nu_n - \nu_n^0\}_{n=0, n \neq m}^\infty$. Since $e^{\nu_m} > 0$, we have

$$P_m : \mathcal{N}^b \rightarrow \mathcal{N}_m^b,$$

where $\mathcal{N}_m^b$ is given by (1.5). Moreover, this mapping is a bijection between $\mathcal{N}^b$ and $\mathcal{N}_m^b$ since ν_m is uniquely reconstructed from $\{\nu_n\}_{n: n \neq m}$ by

$$\nu_m = \log |\dot{w}^*(\lambda_0^*)| + \log \left[2 - b + \sum_{n: n \neq m} \left(2 - \frac{e^{\nu_n}}{|\dot{w}^*(\lambda_n^*)|} \right) \right].$$

Note that Theorem 1.1 yields that the mapping

$$\{\nu_n(q, b) - \nu_n^0\}_0^\infty \mapsto (c^*; \{\mu_n^*\}_0^\infty; \{\nu_n(q, b) - \nu_n^0\}_0^\infty) = \Phi(q, b) \mapsto b = \sum_{n \geq 0} \left(2 - \frac{e^{\nu_n}}{|\dot{w}^*(\lambda_n^*)|} \right)$$

is real-analytic. It is clear that $\partial b / \partial \nu_m = -|\dot{w}(\lambda_m)|^{-1} \neq 0$. Using the Implicit Function Theorem, we obtain that $P_m^{-1} : \mathcal{N}_m^b \rightarrow \mathcal{N}^b$ is real-analytic. Therefore, P_m is a real-analytic isomorphism and the mapping $q \mapsto \{\nu_n(q, b) - \nu_n^0\}_{n=0, n \neq m}^\infty$ is a composition of real-analytic isomorphisms. $\square$

For each $c \in \mathbb{R}$ and $\{\mu_n\}_0^\infty \in \mathcal{M}$, we introduce the function

$$W(\lambda) = W(\lambda; c, \mu_0, \mu_1, \dots) = \cos \sqrt{\lambda} \cdot \prod_{k \geq 0} \frac{\lambda - \lambda_k}{\lambda - \lambda_k^0}, \quad \text{where } \lambda_n = \lambda_n^0 + c + \mu_n, \quad n \geq 0. \quad (4.1)$$

Lemma 4.3. *Let $c \in \mathbb{R}$, $\{\mu_n\}_0^\infty \in \mathcal{M}$ and $n \geq 0$. Then,*

$$\frac{\partial}{\partial \mu_0} |\dot{W}(\lambda_n)|^{-1} > 0, \quad \lim_{\mu_0 \rightarrow -\infty} |\dot{W}(\lambda_n)|^{-1} = 0.$$

Moreover, $\lim_{\mu_0 \rightarrow \lambda_1^0 - \lambda_0^0 + \mu_1} |\dot{W}(\lambda_0)|^{-1} = \lim_{\mu_0 \rightarrow \lambda_1^0 - \lambda_0^0 + \mu_1} |\dot{W}(\lambda_1)|^{-1} = +\infty$.

Remark. Note that the convergence $\mu_0 \rightarrow \lambda_1^0 - \lambda_0^0 + \mu_1$ is equivalent to $\lambda_0 \rightarrow \lambda_1$.

Proof. In order to consider W as a function of μ_0 , we introduce the notation

$$W_\beta(\lambda) = W(\lambda; c, \mu_0 + \beta, \mu_1, \mu_2, \dots) = \frac{\lambda - \lambda_0 - \beta}{\lambda - \lambda_0} \cdot W(\lambda) = \left(1 - \frac{\beta}{\lambda - \lambda_0}\right) W(\lambda).$$

Let $n \geq 1$. Due to $W(\lambda_n) = 0$ and $\dot{W}(\lambda_n) \neq 0$, we have

$$\dot{W}_\beta(\lambda_n) = \left(1 - \frac{\beta}{\lambda_n - \lambda_0}\right) \dot{W}(\lambda_n), \quad \frac{\partial \dot{W}(\lambda_n)}{\partial \mu_0} = \frac{\partial}{\partial \beta} \dot{W}_\beta(\lambda_n) \Big|_{\beta=0} = -\frac{\dot{W}(\lambda_n)}{\lambda_n - \lambda_0}.$$

Hence,

$$\begin{aligned} \frac{\partial}{\partial \mu_0} |\dot{W}(\lambda_n)|^{-1} &= |\dot{W}(\lambda_n)|^{-2} \cdot \frac{|\dot{W}(\lambda_n)|}{\lambda_n - \lambda_0} > 0, \\ \lim_{\mu_0 \rightarrow -\infty} |\dot{W}(\lambda_n)|^{-1} &= \lim_{\beta \rightarrow -\infty} |\dot{W}_\beta(\lambda_n)|^{-1} = \lim_{\beta \rightarrow -\infty} \left|1 - \frac{\beta}{\lambda_n - \lambda_0}\right|^{-1} \cdot |\dot{W}(\lambda_n)|^{-1} = 0. \end{aligned}$$

Moreover, if $n = 1$, then

$$\lim_{\mu_0 \rightarrow \lambda_1^0 - \lambda_0^0 + \mu_1} |\dot{W}(\lambda_1)|^{-1} = \lim_{\beta \rightarrow \lambda_1 - \lambda_0} |\dot{W}_\beta(\lambda_1)|^{-1} = \lim_{\beta \rightarrow \lambda_1 - \lambda_0} \left|1 - \frac{\beta}{\lambda_1 - \lambda_0}\right|^{-1} \cdot |\dot{W}(\lambda_1)|^{-1} = +\infty.$$

Let $n = 0$. We have

$$\dot{W}_\beta(\lambda_0 + \beta) = \frac{W(\lambda_0 + \beta)}{\beta}, \quad \frac{\partial \dot{W}(\lambda_0)}{\partial \mu_0} = \frac{\partial}{\partial \beta} \dot{W}_\beta(\lambda_0 + \beta) \Big|_{\beta=0} = \frac{\ddot{W}(\lambda_0)}{2}.$$

Recall that $\dot{W}(\lambda_0) < 0$. Therefore, the Hadamard factorization yields

$$W(\lambda) = C(\lambda - \lambda_0) \prod_{n \geq 1} \left(1 - \frac{\lambda - \lambda_0}{\lambda_n - \lambda_0}\right), \quad C < 0,$$

and $\ddot{W}(\lambda_0) = -2C \sum_{n \geq 1} (\lambda_n - \lambda_0)^{-1} > 0$. Hence,

$$\frac{\partial}{\partial \mu_0} |\dot{W}(\lambda_0)|^{-1} = -\frac{\partial}{\partial \mu_0} (\dot{W}(\lambda_0))^{-1} = \frac{\ddot{W}(\lambda_0)}{2(\dot{W}(\lambda_0))^2} > 0,$$

$$\lim_{\mu_0 \rightarrow -\infty} |\dot{W}(\lambda_0)|^{-1} = \lim_{\beta \rightarrow -\infty} |W_\beta(\lambda_0 + \beta)|^{-1} = \lim_{\beta \rightarrow -\infty} |\beta| \cdot |W(\lambda_0 + \beta)|^{-1} = 0,$$

and

$$\lim_{\mu_0 \rightarrow \lambda_1^0 - \lambda_0^0 + \mu_1} |\dot{W}(\lambda_0)|^{-1} = \lim_{\beta \rightarrow \lambda_1 - \lambda_0} |W_\beta(\lambda_0 + \beta)|^{-1} = \lim_{\beta \rightarrow \lambda_1 - \lambda_0} |W(\lambda_0 + \beta)|^{-1} \cdot |\beta| = +\infty,$$

since $|W(\lambda_1)| = 0$. □

For each $c \in \mathbb{R}$, $\{\mu_n\}_0^\infty \in \mathcal{M}$ and $\{\nu_n - \nu_n^0\}_0^\infty \in \ell_1^2$, we put

$$B(c, \mu_0, \nu_0, \dots) = \sum_{n \geq 0} \left(2 - \frac{e^{\nu_n}}{|\dot{W}(\lambda_n)|} \right), \quad (4.2)$$

where the function W is given by (4.1). Note that the sum converges due to Theorem 1.1.

Proof of Corollary 1.2. Fix some $b \in \mathbb{R}$ and $m \geq 0$. Due to Theorem 1.1, the mapping

$$q \mapsto \Phi(q, b) = (Q_0 + 2b; \{\mu_n(q, b)\}_0^\infty; \{\nu_n(q, b) - \nu_n^0\}_0^\infty)$$

is a bijection between $L^2(0, 1)$ and the set

$$\mathcal{S}^b = \left\{ (c; \{\mu_n\}_0^\infty; \{\nu_n - \nu_n^0\}_0^\infty) \in \mathbb{R} \times \mathcal{M} \times \ell_1^2 : B(c, \mu_0, \nu_0, \dots) = b \right\}.$$

Define the mapping $P : \mathcal{S}^b \subset \mathbb{R} \times \mathcal{M} \times \ell_1^2 \rightarrow \mathbb{R} \times \mathcal{M}^{(1)} \times \ell_1^2$ by

$$(c; \{\mu_n\}_0^\infty; \{\nu_n - \nu_n^0\}_0^\infty) \mapsto (c; \{\mu_{n+1}\}_0^\infty; \{\nu_n - \nu_n^0\}_0^\infty).$$

Note that $\Phi_b(q) = P(\Phi(q, b))$. In particular, Φ_b is real-analytic as a composition of real-analytic mappings. Due to (4.2) and Lemma 4.3, we have

$$\frac{\partial B(c, \mu_0, \nu_0, \dots)}{\partial \mu_0} < 0, \quad \lim_{\mu_0 \rightarrow -\infty} B(c, \mu_0, \nu_0, \dots) = +\infty, \quad \lim_{\mu_0 \rightarrow \lambda_1^0 - \lambda_0^0 + \mu_1} B(c, \mu_0, \nu_0, \dots) = -\infty.$$

Therefore, for each $(c; \{\mu_{n+1}\}_0^\infty; \{\nu_n - \nu_n^0\}_0^\infty) \in \mathbb{R} \times \mathcal{M}^{(1)} \times \ell_1^2$ there exists unique point μ_0 such that $\mu_0 < \lambda_1^0 - \lambda_0^0 + \mu_1$ and $B(c, \mu_0, \nu_0, \dots) = b$. Thus, the mapping P is a bijection between $\mathcal{S}^b$ and $\mathbb{R} \times \mathcal{M}^{(1)} \times \ell_1^2$. Hence, Φ_b is a bijection between $L^2(0, 1)$ and $\mathbb{R} \times \mathcal{M}^{(1)} \times \ell_1^2$. Using Theorem 1.1, we see that the mapping

$$(c, \mu_0, \nu_0, \dots) \mapsto B(c, \mu_0, \nu_0, \dots)$$

is real-analytic and $\partial B / \partial \mu_0 < 0$ due to Lemma 4.3. Then, the Implicit Function Theorem yields that $P^{-1} : \mathbb{R} \times \mathcal{M}^{(1)} \times \ell_1^2 \rightarrow \mathcal{S}^b$ is real-analytic too. We deduce that $(\Phi^b)^{-1}$ is real-analytic as a composition of real-analytic mappings. □

5 Appendix. The case of general boundary conditions

Consider the Sturm-Liouville problem

$$\begin{aligned} -\psi'' + q(x)\psi &= \lambda\psi, \quad x \in [0, 1], \quad q \in L^2(0, 1), \\ \psi'(0) - a\psi(0) &= 0, \quad \psi'(1) + b\psi(1) = 0, \quad a, b \in \mathbb{R}. \end{aligned}$$

Denote by $\sigma_n(q, a, b)$ the eigenvalues of this problem. It is well-known that

$$\sigma_n(q, a, b) = \sigma_n^0 + Q_0 + 2a + 2b + \tau_n(q, a, b), \quad \text{where} \quad Q_0 = \int_0^1 q(t)dt, \quad \{\tau_n\}_0^\infty \in \ell^2 \quad (5.1)$$

and $\sigma_n^0 = \pi^2 n^2$, $n \geq 0$. Note the $\sigma_n(q, a, b)$ are the roots of the Wronskian

$$w(\lambda, q, a, b) = (\vartheta' + a\varphi' + b(\vartheta + a\varphi))(1, \lambda, q).$$

Following [IT], we introduce the norming constants

$$\varkappa_n(q, a, b) = \log[(-1)^n(\vartheta + a\varphi)(1, \sigma_n, q)] = \log \left| \frac{\psi_n(1, q, a, b)}{\psi_n(0, q, a, b)} \right|,$$

where ψ_n is the n -th normalized eigenfunction such that $\psi_n(0) > 0$.

Theorem 5.1 (Isaacson, Trubowitz [IT]). *The mapping*

$$\Psi : (q; a; b) \mapsto (Q_0 + 2a + 2b; \{\tau_n(q, a, b)\}_0^\infty; \{\varkappa_n(q, a, b)\}_0^\infty)$$

is a real-analytic isomorphism between $L^2(0, 1) \times \mathbb{R}^2$ and $\mathbb{R} \times \mathcal{T} \times \ell_1^2$, where

$$\mathcal{T} = \{\{\tau_n\}_0^\infty \in \ell^2 : \sigma_0^0 + \tau_0 < \sigma_1^0 + \tau_1 < \dots\} \subset \ell^2.$$

Theorem 5.1 and Theorem 1.1 are similar. In order to consider the case of fixed $a, b \in \mathbb{R}$, we need some modification of identity (1.4).

Proposition 5.2. *For each $(q; a; b) \in L^2(0, 1) \times \mathbb{R}^2$ the following identities are fulfilled:*

$$-1 + \sum_{n \geq 0} \left(2 - \frac{e^{\varkappa_n}}{|\dot{w}(\sigma_n)|} \right) = b; \quad -1 + \sum_{n \geq 0} \left(2 - \frac{e^{-\varkappa_n}}{|\dot{w}(\sigma_n)|} \right) = a, \quad (5.2)$$

where $w(\lambda) = w(\lambda, q, a, b) = -\sqrt{\lambda} \sin \sqrt{\lambda} \cdot \prod_{n \geq 0} \frac{\lambda - \sigma_n}{\lambda - \sigma_n^0}$ and $\sigma_n = \sigma_n(q, a, b)$, $\varkappa_n = \varkappa_n(q, a, b)$.

Remark. Note that identities (5.2) and Theorem 5.1 directly yield the following well-known result:

Fix some constant $a = b$ and $q \in L_{\text{even}}^2(0, 1)$, i.e. $q(x) = q(1 - x)$, $x \in [0, 1]$.

If $p \in L^2(0, 1)$ is such that $\sigma_n(p, a, a) = \sigma_n(q, a, a)$ for all $n \geq 0$, then $p = q$.

Indeed, in this case we have $w(\lambda, p, a, a) = w(\lambda, q, a, a)$ and $\varkappa_n(p, a, a) = 0$ for all $n \geq 0$. Summing identities (5.2) for the potential p and subtracting corresponding ones for the potential q , we deduce that

$$\sum_{n \geq 0} \frac{\cosh \varkappa_n(p, a, a) - 1}{|\dot{w}(\sigma_n)|} = 0.$$

This yields $\varkappa_n(p, a, a) = 0$ for all $n \geq 0$ since $\cosh \varkappa_n \geq 1$. Thus, Theorem 5.1 gives $p = q$.

Let $a, b \in \mathbb{R}$ be fixed. Then, due to identities (5.2), it is possible to reconstruct $(\sigma_0, \varkappa_0)$ from the other spectral data $\{\sigma_n\}_1^\infty, \{\varkappa_n\}_1^\infty$. More precisely, we have

Proposition 5.3. *For any fixed $a, b \in \mathbb{R}$ the mapping*

$$\Psi_{a,b} : q \mapsto (Q_0 + 2a + 2b; \{\tau_{n+1}(q, a, b)\}_0^\infty; \{\varkappa_{n+1}(q, a, b)\}_0^\infty)$$

is a real-analytic isomorphism between $L^2(0, 1)$ and $\mathcal{T}^{(1)} \times \ell_1^2$, where

$$\mathcal{T}^{(1)} = \{\{\tau_{n+1}\}_0^\infty \in \ell^2 : \sigma_1^0 + \tau_1 < \sigma_2^0 + \tau_2 < \dots\} \subset \ell^2.$$

Proof of Proposition 5.2. We prove the first identity in (5.2), the proof of the second is similar. Consider the meromorphic function

$$f(\lambda) = \frac{(\vartheta + a\varphi)(1, \lambda, q)}{w(\lambda, q, a, b)} = \frac{\vartheta + a\varphi}{(\vartheta' + a\varphi') + b(\vartheta + a\varphi)}(1, \lambda, q), \quad \lambda \in \mathbb{C}.$$

All roots $\sigma_n = \sigma_n(q, a, b)$ of $w(\lambda)$ are simple. Therefore, definition (5.1) yields

$$\operatorname{res}_{\lambda=\sigma_n} f(\lambda) = \frac{(\vartheta + a\varphi)(1, \lambda, q)}{\dot{w}(\sigma_n)} = \frac{(-1)^n e^{\varkappa_n}}{\dot{w}(\sigma_n)} = -\frac{e^{\varkappa_n}}{|\dot{w}(\sigma_n)|}. \quad (5.3)$$

Put $|\lambda| = \pi^2(m + \frac{1}{2})^2 \rightarrow \infty$. Then, due to asymptotics (2.2)-(2.5), we have

$$\begin{aligned} f(\lambda) &= \frac{\cos \sqrt{\lambda} + \lambda^{-1/2} \sin \sqrt{\lambda} \cdot (\frac{1}{2}Q_0 + a) + o(\lambda^{-1/2}e^{|\operatorname{Im} \sqrt{\lambda}|})}{-\lambda^{1/2} \sin \sqrt{\lambda} + \cos \sqrt{\lambda} \cdot (\frac{1}{2}Q_0 + a + b) + o(e^{|\operatorname{Im} \sqrt{\lambda}|})} \\ &= -\lambda^{-\frac{1}{2}} \cot \lambda - \lambda^{-1} \cdot (\frac{1}{2}Q_0 + a) - \lambda^{-1} \cot^2 \lambda \cdot (\frac{1}{2}Q_0 + a + b) + o(\lambda^{-1}). \end{aligned}$$

Let $f_0(\lambda) = -\lambda^{-\frac{1}{2}} \cot \sqrt{\lambda}$. Applying the Cauchy Theorem in the disk $|\lambda| < \pi^2(m + \frac{1}{2})^2$, we obtain

$$\sum_{n=0}^m \left(\operatorname{res}_{\lambda=\sigma_n} f(\lambda) - \operatorname{res}_{\lambda=\sigma_n^0} f_0(\lambda) \right) = -(\frac{1}{2}Q_0 + a) - (\frac{1}{2}Q_0 + a + b) \cdot \sum_{n=0}^m \operatorname{res}_{\lambda=\sigma_n^0} (f_0(\lambda))^2. \quad (5.4)$$

Note that

$$f_0(\lambda) = -\frac{1}{\sqrt{\lambda} \tan \sqrt{\lambda}} = -\frac{1}{\lambda} \left(1 - \frac{\lambda}{3} + O(\lambda^2) \right), \quad \lambda \rightarrow \sigma_0 = 0,$$

and

$$f_0(\lambda) = -\frac{2}{\lambda - \sigma_n^0} \left(1 - \frac{\lambda - \sigma_n^0}{4\sigma_n^0} + O((\lambda - \sigma_n^0)^2) \right), \quad \lambda \rightarrow \sigma_n^0, \quad n \geq 1.$$

Hence,

$$\operatorname{res}_{\lambda=\sigma_0^0} f_0(\lambda) = -1, \quad \operatorname{res}_{\lambda=\sigma_n^0} f_0(\lambda) = -2, \quad n \geq 1,$$

$$\operatorname{res}_{\lambda=\sigma_0^0} (f_0(\lambda))^2 = -\frac{2}{3}, \quad \operatorname{res}_{\lambda=\sigma_n^0} (f_0(\lambda))^2 = -\frac{2}{\sigma_n^0}, \quad n \geq 1,$$

and

$$\sum_{n \geq 0} \operatorname{res}_{\lambda=\sigma_n^0} (f_0(\lambda))^2 = -\frac{2}{3} - \frac{2}{\pi^2} \sum_{n \geq 0} \frac{1}{n^2} = -\frac{2}{3} - \frac{2}{\pi^2} \cdot \frac{\pi^2}{6} = -1.$$

Substituting these identities and (5.3) into (5.4), we obtain (5.2) as $m \rightarrow \infty$. $\square$

For each $c \in \mathbb{R}$, $\{\tau_n\}_0^\infty \in \mathcal{T}$ and $\{\varkappa_n\}_0^\infty \in \ell_1^2$ we put

$$W(\lambda) = W(\lambda; c, \tau_0, \tau_1, \dots) = -\sqrt{\lambda} \cos \sqrt{\lambda} \cdot \prod_{k \geq 0} \frac{\lambda - \sigma_k}{\lambda - \sigma_k^0}, \quad \text{where } \sigma_k = \sigma_k^0 + c + \tau_k, \quad k \geq 0.$$

$$F(c, \tau_0, \tau_1, \dots) = \frac{1}{|\dot{W}(\sigma_0)|}, \quad G_\pm(c, \tau_0, \tau_1, \varkappa_1, \tau_2, \varkappa_2, \dots) = -1 + \sum_{n \geq 1} \left(\frac{e^{\pm \varkappa_n}}{|\dot{W}(\sigma_n)|} - 2 \right). \quad (5.5)$$

Note that the sum in the definition of $G_\pm$ converges due to Theorem 5.1 and Proposition 5.2. Using these notations, we rewrite the identities (5.2) in the following form:

$$e^{-\varkappa_0} F + G_- = -a, \quad e^{\varkappa_0} F + G_+ = -b. \quad (5.6)$$

Lemma 5.4. (i) Let $c \in \mathbb{R}$, $\{\tau_n\}_0^\infty \in \mathcal{T}$ and $\{\varkappa_n\}_0^\infty \in \ell_1^2$. Then,

$$\frac{\partial F}{\partial \tau_0} > 0, \quad \lim_{\tau_0 \rightarrow -\infty} F = 0, \quad \lim_{\tau_0 \rightarrow \sigma_1^0 - \sigma_0^0 + \tau_1} F = +\infty; \quad (5.7)$$

$$\frac{\partial G_\pm}{\partial \tau_0} > 0, \quad \lim_{\tau_0 \rightarrow -\infty} G_\pm = -\infty, \quad \lim_{\tau_0 \rightarrow \sigma_1^0 - \sigma_0^0 + \tau_1} G_\pm = +\infty. \quad (5.8)$$

(ii) For each $a, b, c \in \mathbb{R}$, $\{\tau_{n+1}\}_0^\infty \in \mathcal{T}^{(1)}$, $\{\varkappa_{n+1}\}_0^\infty \in \ell_1^2$ there exist unique $\tau_0 < \sigma_1^0 - \sigma_0^0 + \tau_1$ and $\varkappa_0 \in \mathbb{R}$ such that identities (5.6) (or, equivalently, (5.2)) hold true.

Proof. (i) The proof repeats the proof of Lemma 4.3.

(ii) Note that identities (5.6) are equivalent to

$$F^2 = (a + G_-)(b + G_+), \quad e^{\varkappa_0} = -\frac{F}{a + G_-} = -\frac{b + G_+}{F}.$$

We will consider F and $G_\pm$ as functions of τ_0 . It follows from (5.7) that the function $F(\tau_0)$ is positive for all $\tau_0 < \sigma_1^0 - \sigma_0^0 + \tau_1$. Hence, $a + G_-(\tau_0) < 0$ and $b + G_+(\tau_0) < 0$. Due to (5.8), there exist unique points $\tau_\pm^* < \sigma_1^0 - \sigma_0^0 + \tau_1$ such that $a + G_-(\tau_-^*) = b + G_+(\tau_+^*) = 0$. Since the functions $G_\pm(\tau_0)$ are increasing, we have $\tau_0 < \tau_\pm^*$. Put $\tau^* = \min(\tau_-^*, \tau_+^*)$ and

$$G(\tau_0) = (a + G_-(\tau_0)) \cdot (b + G_+(\tau_0)), \quad \tau_0 \leq \tau_*.$$

Using (5.8), it is easy to see that

$$G'(\tau_0) < 0, \quad \lim_{\tau_0 \rightarrow -\infty} G(\tau_0) = +\infty, \quad G(\tau_*) = 0; \quad (F^2)'(\tau_0) > 0, \quad \lim_{\tau_0 \rightarrow -\infty} (F(\tau_0))^2 = 0.$$

Therefore, the equation $(F(\tau_0))^2 = G(\tau_0)$ has the unique solution $\tau_0 \in (-\infty, \tau_*)$ and $\varkappa_0 = \log |F(\tau_0)/(a + G_-(\tau_0))| = \log |(b + G_+(\tau_0))/F(\tau_0)|$ is uniquely determined by τ_0 . $\square$

Proof of Proposition 5.3. Fix some $a, b \in \mathbb{R}$ and $m \geq 0$. Due to Theorem 5.1 and Proposition 5.2, the mapping $q \mapsto \Psi(q, a, b) = (Q_0 + 2a + 2b; \{\tau_n(q, a, b)\}_0^\infty; \{\varkappa_n(q, a, b)\}_0^\infty)$ is a bijection between $L^2(0, 1)$ and the set

$$\mathcal{S}^{a,b} = \left\{ (c; \{\tau_n\}_0^\infty; \{\varkappa_n\}_0^\infty) \in \mathbb{R} \times \mathcal{T} \times \ell_1^2 : e^{-\varkappa_0} F + G_- = -a \text{ and } e^{\varkappa_0} F + G_+ = -b \right\},$$

where F and $G_\pm$ are given by (5.5).

Define the mapping $P : \mathcal{S}^{a,b} \subset \mathbb{R} \times \mathcal{T} \times \ell_1^2 \rightarrow \mathbb{R} \times \mathcal{T}^{(1)} \times \ell_1^2$ by

$$(c; \{\tau_n\}_0^\infty; \{\varkappa_n\}_0^\infty) \mapsto (c; \{\tau_{n+1}\}_0^\infty; \{\varkappa_n\}_0^\infty).$$

Note that $\Psi_{a,b}(q) = P(\Psi(q, a, b))$. In particular, $\Psi_{a,b}$ is real-analytic as a composition of real-analytic mappings. Due to Lemma 5.4 (ii), for each $(c; \{\tau_{n+1}\}_0^\infty; \{\varkappa_n\}_0^\infty) \in \mathbb{R} \times \mathcal{M}^{(1)} \times \ell_1^2$ there exist unique $\tau_0 < \sigma_1^0 - \sigma_0^0 + \tau_1$ and $\varkappa_0 \in \mathbb{R}$ such that both equations $e^{-\varkappa_0}F + G_- = -a$ and $e^{\varkappa_0}F + G_+ = -b$ hold true. Therefore, the mapping P is a bijection between $\mathcal{S}^{a,b}$ and $\mathbb{R} \times \mathcal{M}^{(1)} \times \ell_1^2$. Thus, $\Psi_{a,b}$ is a bijection between $L^2(0, 1)$ and $\mathbb{R} \times \mathcal{M}^{(1)} \times \ell_1^2$.

In order to prove that $\Psi_{a,b}^{-1}$ is real-analytic, we deduce from Theorem 5.1 that the mapping

$$(c; \{\tau_n\}_0^\infty; \{\varkappa_n\}_0^\infty) \mapsto (-a; -b) = (e^{-\varkappa_0}F + G_-; e^{\varkappa_0}F + G_+)$$

is real-analytic. Lemma 5.4 (i) gives $\frac{\partial a}{\partial \tau_0} < 0$, $\frac{\partial b}{\partial \tau_0} < 0$, $\frac{\partial a}{\partial \varkappa_0} = e^{-\varkappa_0}F > 0$, $\frac{\partial b}{\partial \varkappa_0} = e^{\varkappa_0}F < 0$. Hence,

$$\det \begin{pmatrix} \partial a / \partial \tau_0 & \partial b / \partial \tau_0 \\ \partial a / \partial \varkappa_0 & \partial b / \partial \varkappa_0 \end{pmatrix} > 0.$$

Then, the Implicit Function Theorem yields that $P^{-1} : \mathbb{R} \times \mathcal{T}^{(1)} \times \ell_1^2 \rightarrow \mathcal{S}^{a,b}$ is real-analytic. We obtain that $(\Phi^b)^{-1}$ is real-analytic as a composition of real-analytic mappings. $\square$

References

- [CK] Chelkak D., Korotyaev E.: The inverse problem for perturbed harmonic oscillator on the half-line. Preprint 2005
- [CKK] Chelkak D., Kargaev P., Korotyaev E.: Inverse Problem for Harmonic Oscillator Perturbed by Potential, Characterization. *Comm. Math. Phys.* **249**(1), 133–196 (2004)
- [DT] Dahlberg B., Trubowitz E.: The Inverse Sturm-Liouville Problem III. *Comm. Pure Appl. Math.* **37**, 255–267 (1984)
- [IT] Isaacson E., Trubowitz E.: The Inverse Sturm-Liouville Problem I. *Comm. Pure Appl. Math.* **36**, 767–783 (1983)
- [IMT] Isaacson E., McKean H.P., Trubowitz E.: The Inverse Sturm-Liouville Problem II. *Comm. Pure Appl. Math.* **37**, 1–11 (1984)
- [JL] Jodeit, M.J., Levitan, B. M.: The isospectrality problem for the classical Sturm-Liouville equation. *Adv. Differential Equations* **2**(2), 297–318 (1997).
- [L] Levitan, B.: *Inverse Sturm-Liouville problems*. Utrecht: VNU Science Press, 1987.
- [M] Marchenko, V.: *Sturm-Liouville operator and applications*. Basel: Birkhäuser, 1986.
- [PT] Pöschel P., Trubowitz E.: *Inverse Spectral Theory*. Boston: Academic Press, 1987.